



Please read carefully the use of information

MANACIL

Drug solution for the treatment of ornamental fish in fresh water aquariums

manaus-aquarium

Rüchnerstrasse 73
D-64319 Pfungstadt / Germany
☎ +49 6157 1587230
✉ info@manaus-aquarium.com
www.manaus-aquarium.com

Pharmaceutical for animals: (Fresh water ornamental fishes)

Pharmaceutical contractor:

solution of active pharmaceutical ingredients to load water in aquariums.

Consistency of effective components mg pro ml:

Brilliantgrün 48, 4,4'-Benzyliden-Bis(n,n-Dimethylaniline) 9.6, Acriflavin HCL 4, Matanigelb 0.05

Pharmaceutical Entrepreneurs:

Manaus-Aquarium, Rüchnerstrasse 73, D-64319 Pfungstadt

Field of application: Very effectively **MANACIL** combats (Ichthyophthirius multifiliis) as well as monocellular parasites of the skin of fishes. It also combats Chilodonella, Ichthyobodo necator, Trichodina etc..

Symptoms:

Indications of affection are white dots upon the skin and fins (Ichthyo). All other pathogens mentioned above are shown as grey or white coverings on the skin of the fishes. In general ill fishes refuse feeding. They "clip" their fins together, the behaviour gets jumpy and they try to rub on objects in the aquarium. It is important, to medicate the fishes right after diagnosing the symptoms, in order to prevent weakening the fishes by the viruses mentioned above. It is recommended to diagnose by microscope – if possible.

Preventive measures:

Don't use **MANACIL** in combination with other medicine. If your fishes were treated with other medicine before, it is necessary to change the water before treatment.

Pay attention to the quality of the water and reduce the organic contamination by chemical decomposition products by changing the water of the aquarium.

Strongly polluted filters should be cleaned before the therapy. Filter and air ducts (?Ausströmer?) for oxygen supply should always stay in progress! Ultraviolet lamps have to be switched off during the therapy.

Do not make use of active filter media (for example active carbon, Phosphatentferner etc.) in filters during treatment, to avoid interactions with bonded interactive substances.

Remove the ion exchanger (for example for removal of nitrate etc.) during the therapy with **MANACIL**.

In case of complications stop the treatment with **MANACIL** and change the water – as usual when medicating ornamental fishes and inform the producer.

Dosing and application:

Use the measuring cup enclosed to the package to dose *MANACIL* and spread the medicine on the surface of the water.

1st day: 1 ml per 40 litres water

3rd day: 1 ml per 40 litres water

In case of an intense affection with parasites you may use again 1ml per 40 litres on the 5th day.

Do not overdose *MANACIL* – fishes weakened by illness may react sensitively.

Contraindication and adverse reactions:

MANACIL is a new product and well biocompatible.

Adverse reactions are not known.

Keepers of ornamental fishes should inform the veterinarian or pharmacist and the producer about any unknown adverse reaction.

Important warning:

Keep out of reach of children!

Don't make use of the medicine on animals that are kept to produce foodstuffs.

Keep it in the original package and defend it from light and heat.

Store it frost-protected.

Do not make use of it after the expiry date is exceeded.

MANACIL can contribute coloring